


Aspirin


aspirin 325mg

PART NO.: 20-155

Packets



Side 1



Side 2



Side 3



Side 4

Aspirin

aspirin 325mg

PART NO.: 90014

Packets

PhysiciansCARE[®] 2 Tablets

ASPIRIN

Active ingredients (in each tablet)

Purpose

Aspirin (NSAID) 325 mg Pain reliever / fever reducer
nonsteroidal anti-inflammatory drug

Uses

Temporarily relieves

headache ■ muscular aches ■ minor arthritis pain ■ backache
pain and fever of colds ■ toothache ■ menstrual cramps

Temporarily reduces fever

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling ■ shock ■ asthma (wheezing) ■ anaphylaxis ■ anaphylactic shock. If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinner (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcohol drinks every day while using this product ■ take more or for a longer time than directed

Package Not Child-Resistant

Tamper-evident sealed packets

Do not use any opened or torn packets

Store at 59-86°F (15-30°C) in a dry place

Avoid excessive heat above 104°F (40°C)

Manufactured for Acme United Corp

55 Walls Drive, Fairfield, CT 06424 • 800.835.2263

Warnings (continued)

Do not use ■ if you are allergic to aspirin or to any other pain reliever/fever reducer

Ask a doctor before use if ■ stomach bleeding warning applies to you ■ you have a history of stomach problems such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic ■ you have asthma

Ask a doctor or pharmacist before use if you are ■ taking prescription drugs for gout, diabetes or arthritis

When using this product ■ take with food or milk if stomach upset occurs

Stop use and ask a doctor if ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better

■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ any new symptoms appear ■ if ringing in the ears or loss of hearing occurs ■ redness or swelling is present in the painful area ■ an allergic reaction occurs and seek medical help right away

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

Drink a full glass of water with each dose

Adults and children (12 years and older) Take 1 or 2 tablets with water every 4 hours as needed. Do not take more than 12 tablets in 24 hours, or as directed by a doctor.

Children under 12 years Do not give to children under 12 years of age.

SEE CARTON FOR COMPLETE PRODUCT INFORMATION

90014

PhysiciansCARE[®]

ASPIRIN

Pain Reliever and Fever Reducer (NSAID)

THIS PACKAGE FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN.

COATED TABLETS

Tamper-Evident Packets of 2 Tablets
Do not use if packet is open or torn.

Aspirin (NSAID) 325 mg
100 Tablets (50 Packets, 2 tablets each)

Side 1

90014

Drug Facts

Active ingredient (in each tablet)

Purpose

Aspirin (NSAID) 325 mg Pain reliever / fever reducer
nonsteroidal anti-inflammatory drug

Uses

Temporarily relieves:

headache ■ minor arthritis pain ■ pain & fever of colds
backache ■ menstrual cramps ■ toothache ■ muscular aches

Temporarily reduces fever

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling ■ shock ■ asthma (wheezing) ■ anaphylaxis ■ anaphylactic shock. If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinner (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcohol drinks every day while using this product ■ take more or for a longer time than directed

Do not use

■ if you are allergic to aspirin or any other pain reliever/fever reducer

Ask a doctor before use if

stomach bleeding warning applies to you
you have asthma
you have a history of stomach problems such as heartburn
you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
you are taking a diuretic

Ask a doctor or pharmacist before use if you are

■ if you are taking prescription drugs for gout, diabetes or arthritis

When using this product

■ take with food or milk if stomach upset occurs

Drug Facts (continued)

Stop use and ask a doctor if

you experience any of the following signs of stomach bleeding:
■ feel faint
■ vomit blood
■ have bloody or black stools
■ have stomach pain that does not get better
■ pain gets worse or lasts more than 10 days
■ fever gets worse or lasts more than 3 days
■ if ringing in the ears or loss of hearing occurs
■ redness or swelling is present in the painful area
■ any new symptoms appear
■ an allergic reaction occurs. Seek medical help right away.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

Drink a full glass of water with each dose

Adults and children (12 years and older) Take 1 or 2 tablets with water every 4 hours as needed. Do not take more than 12 tablets in 24 hours, or as directed by a doctor.

Children under 12 years Do not give to children under 12 years of age.

Other information

read all product information before using
store at room temperature 59-86°F (15-30°C)
avoid excessive heat and humidity
do not use any opened or torn packets

Inactive ingredients

corn starch, hypromellose, polyethylene glycol, propylene glycol

Questions? 1-800-835-2263

Contact Us 1-800-835-2263 www.FirstAidOnly.com

Side 2

90014

PhysiciansCARE[®]

ASPIRIN

Pain Reliever and Fever Reducer (NSAID)

THIS PACKAGE FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN.

COATED TABLETS

Tamper-Evident Packets of 2 Tablets
Do not use if packet is open or torn.

Aspirin (NSAID) 325 mg
100 Tablets (50 Packets, 2 tablets each)

Side 3

Also Available from PhysiciansCare[®] in individual dose packets:

PhysiciansCare[®] Non-Aspirin (Acetaminophen)

PhysiciansCare[®] Ibuprofen

PhysiciansCare[®] Cold & Cough

PhysiciansCare[®] Allergy

PhysiciansCare[®] Sinus

PhysiciansCare[®] Antacid


Side 4

Aspirin

aspirin 325mg

PART NO.: 54034

Packets



Side 1



Side 2



Side 3





Side 4

Aspirin

aspirin 325mg

PART NO.: J412

Packets



Side 1

Side 2

Side 3

Side 4

Acetaminophen

Acetaminophen 500 mg

PART NO.: 54036

Packets

PhysiciansCARE 2 Tablets

**EXTRA STRENGTH
NON-ASPIRIN**

Active ingredient (in each tablet) **Purpose**
Acetaminophen 500 mg.....Pain reliever/fever reducer

Uses
For the temporary relief of minor aches and pains associated with:
■ headache ■ muscular aches ■ minor arthritis pain
■ toothache ■ common cold ■ menstrual cramps
For the reduction of fever.

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:
■ more than 4,000 mg in 24 hours, which is the maximum daily amount
■ with other drugs containing acetaminophen
■ 3 or more alcoholic drinks every day while using this product

Package Not Child-Resistant
Tamper-evident sealed packets
Do not use any opened or torn packets

Store at 59°-86°F (15°-30°C)
Manufactured for Acme United Corp
55 Walls Drive, Fairfield, CT 06424 • 800-835-2263

Warnings (continued)
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
■ skin reddening ■ blisters ■ rash If a skin reaction occurs, stop use and seek medical help right away.

Do not use
■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
■ for more than 10 days for pain unless directed by a doctor
■ for more than 3 days for fever unless directed by a doctor

Ask a doctor before use if you have
■ liver disease
Ask a doctor or pharmacist before use if
■ you are taking the blood thinning drug warfarin

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.


Directions ■ do not use more than directed
Adults and children: (12 years and older) Take 2 tablets every 6 hours while symptoms last. Do not take more than 8 tablets in 24 hours unless directed by a doctor.
Children under 12 years: Do not give this adult strength product to children under 12 years of age; this will provide more than the recommended dose (overdose) and may cause liver damage.

SEE CARTON FOR COMPLETE PRODUCT INFORMATION

Lot 7102 Exp. 08-2023



Side 1



Side 2



Side 3




Side 4

Acetaminophen


Acetaminophen 500 mg

PART NO.: 90016

Packets




Side 1



Side 2



Side 3



Side 4

Acetaminophen

Acetaminophen 500 mg

PART NO.: 40800

Packets

PhysiciansCARE® 2 Tablets

EXTRA STRENGTH

NON-ASPIRIN

Active ingredient (in each tablet) Purpose

Acetaminophen 500 mg Pain reliever/fever reducer

Uses

For the temporary relief of minor aches and pains associated with:

- headache
- muscular aches
- minor arthritis pain
- toothache
- common cold
- menstrual cramps

For the reduction of fever.

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Package Not Child-Resistant

Tamper-evident sealed packets

Do not use any opened or torn packets

Store at 59°-86°F (15°-30°C)

Manufactured for Acme United Corp

55 Walls Drive, Fairfield, CT 06424 • 800.835.2263

Warnings (continued)

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- for more than 10 days for pain unless directed by a doctor
- for more than 3 days for fever unless directed by a doctor

Ask a doctor before use if you have

- liver disease
- are taking the blood thinning drug warfarin

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not use more than directed

Adults and children (12 years and older) Take 2 tablets every 6 hours while symptoms last. Do not take more than 8 tablets in 24 hours unless directed by a doctor.


Children under 12 years: Do not give this adult strength product to children under 12 years of age; this will provide more than the recommended dose (overdose) and may cause liver damage.

SEE CARTON FOR COMPLETE PRODUCT INFORMATION

Lot 7102 Exp. 08-2023




Side 1



Side 2



Side 3



Side 4

Acetaminophen

Acetaminophen 500 mg

PART NO.: J420

Packets

PhysiciansCARE 2 Tablets

EXTRA STRENGTH

NON-ASPIRIN

Active ingredient (in each tablet) **Purpose**
Acetaminophen 500 mg Pain reliever/fever reducer

Uses
For the temporary relief of minor aches and pains associated with:
■ headache ■ muscular aches ■ minor arthritis pain
■ toothache ■ common cold ■ menstrual cramps
For the reduction of fever.

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:
■ more than 4,000 mg in 24 hours, which is the maximum daily amount
■ with other drugs containing acetaminophen
■ 3 or more alcoholic drinks every day while using this product

Package Not Child-Resistant
Tamper-evident sealed packets
Do not use any opened or torn packets

Store at 59°-86°F (15°-30°C)
Manufactured for Acme United Corp
55 Walls Drive, Fairfield, CT 06424 • 800.835.2263

Warnings (continued)
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
■ skin reddening ■ blisters ■ rash If a skin reaction occurs, stop use and seek medical help right away.

Do not use
■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
■ for more than 10 days for pain unless directed by a doctor
■ for more than 3 days for fever unless directed by a doctor

Ask a doctor before use if you have
■ liver disease
■ you are taking the blood thinning drug warfarin

Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions ■ do not use more than directed
Adults and children (12 years and older) Take 2 tablets every 6 hours while symptoms last. Do not take more than 8 tablets in 24 hours unless directed by a doctor.
Children under 12 years: Do not give this adult strength product to children under 12 years of age; this will provide more than the recommended dose (overdose) and may cause liver damage.
SEE CARTON FOR COMPLETE PRODUCT INFORMATION




Side 1



Side 2



Side 3



Side 4

Acetaminophen

Acetaminophen 500 mg

PART NO.: 91272

Packets

PhysiciansCARE[®] 2 Tablets

EXTRA STRENGTH

NON-ASPIRIN

Active ingredient (in each tablet) Purpose

Acetaminophen 500 mg Pain reliever/fever reducer

Uses

For the temporary relief of minor aches and pains associated with:

- headache
- muscular aches
- minor arthritis pain
- toothache
- common cold
- menstrual cramps

For the reduction of fever.

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Package Not Child-Resistant

Tamper-evident sealed packets

Do not use any opened or torn packets

Store at 59°-86°F (15°-30°C)

Manufactured for Acme United Corp

55 Walls Drive, Fairfield, CT 06424 • 800-835-2263

Warnings (continued)

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use:

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- for more than 10 days for pain unless directed by a doctor
- for more than 3 days for fever unless directed by a doctor

Ask a doctor before use if you have:

- liver disease

Ask a doctor or pharmacist before use if:

- you are taking the blood thinning drug warfarin

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not use more than directed

Adults and children (12 years and older): Take 2 tablets every 6 hours while symptoms last. Do not take more than 8 tablets in 24 hours unless directed by a doctor.


Children under 12 years: Do not give this brand to children under 12 years of age. This will provide more than the recommended dose (overdose) and may cause liver damage.

SEE CARTON FOR COMPLETE PRODUCT INFORMATION

Lot 7102 Exp. 08-2023




Side 1



Side 2



Side 3




Side 4

Pain Relief


Acetaminophen 250 mg
Aspirin 250 mg
Caffeine 65 mg

PART NO.: 90316

Packets




Side 1



Side 2



Side 3




Side 4

Pain Relief

Acetaminophen 250 mg
Aspirin 250 mg
Caffeine 65 mg

PART NO.: 90317

Packets



Side 1



Side 2



Side 3




Side 4

Ibuprofen


Ibuprofen 200 mg

PART NO.: 90015

Packets




Side 1



Side 2



Side 3




Side 4

Ibuprofen

Ibuprofen 200 mg

PART NO.: 90109

Packets




Side 1



Side 2



Side 3



Side 4

Ibuprofen

Ibuprofen 200 mg

PART NO.: J432

Packets

PhysiciansCARE[®] 2 Tablets

IBUPROFEN

Active ingredient (in each tablet) Purpose

Ibuprofen (NSAID) 200mg Pain reliever/fever reducer

*nonsteroidal anti-inflammatory drug

Uses

Temporarily relieves minor aches and pains associated with

• headache • toothache • backache

• menstrual cramps • common cold

• muscular aches • minor arthritis pain

Temporarily reduces fever.

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

• hives • skin reddening • facial swelling • rash

• asthma (wheezing) • blisters • shock

If an allergic reaction occurs, stop use and seek medical help right away.

Package Not Child-Resistant

Tamper-evident sealed packets

Do not use if packet is open or torn.

Store at 68-77°F (20-25°C)

Avoid excessive heat, above 104°F (40°C)

Manufactured for Acme United Corp.

55 Walls Drive, Fairfield, CT 06424 • 800.835.2263

Warnings (continued)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) • have 3 or more alcoholic drinks every day while using this product • take more or for a longer time than directed.

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless specifically directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions • do not use more than directed • the smallest effective dose should be used • do not take longer than 10 days unless directed by a doctor (see warnings)

Adults and children (12 years and older): Take 1 tablet every 4 to 6 hours while symptoms persist. If pain or fever does not respond to 1 tablet, 2 tablets may be used. Do not exceed 6 tablets in 24 hours, unless directed by a doctor.

Children under 12 years: Do not give to children under 12 years of age.

SEE CARTON FOR COMPLETE PRODUCT INFORMATION




Side 1



Side 2



Side 3




Side 4

Ibuprofen


Ibuprofen 200 mg

PART NO.: 91277

Packets




Side 1



Side 2



Side 3



Side 4

Medication Station/Mutli-Pack

Ibuprofen 200 mg, aspirin 325mg,
Acetaminophen 500 mg,
calcium carbonate 420 mg

PART NO.: 90780

Packets



Side 1



Side 2

Medication Station/Mutli-Pack

Ibuprofen 200 mg, aspirin 325mg,
Acetaminophen 500 mg,
calcium carbonate 420 mg

PART NO.: 91273

Packets



Side 1



Side 2